

# Harvard Mental Health Letter

VOLUME 28 • NUMBER 1 | JULY 2011

## How addiction hijacks the brain

*Desire initiates the process, but learning sustains it.*

**T**he word "addiction" is derived from a Latin term for "enslaved by" or "bound to." Anyone who has struggled to overcome an addiction—or has tried to help someone else to do so—understands why.

Addiction exerts a long and powerful influence on the brain that manifests in three distinct ways: craving for the object of addiction, loss of control over its use, and continuing involvement with it despite adverse consequences. While overcoming addiction is possible, the process is often long, slow, and complicated. It took years for researchers and policymakers to arrive at this understanding.

In the 1930s, when researchers first began to investigate what caused addictive behavior, they believed that people who developed addictions were somehow morally flawed or lacking in willpower. Overcoming addiction, they thought, involved punishing miscreants or, alternately, encouraging them to muster the will to break a habit.

The scientific consensus has changed since then. Today we recognize addiction as a chronic disease that changes both brain structure and function. Just as cardiovascular disease damages the heart and diabetes impairs the pancreas, addiction hijacks the brain. Recovery from addiction involves willpower, certainly, but it is not enough to "just say no"—as the 1980s slogan suggested. Instead, people typically use multiple strategies—including psychotherapy, medication, and self-care—as they try to break the grip of an addiction.

Another shift in thinking about addiction has occurred as well. For many years, experts believed that only alcohol and powerful drugs could cause addiction. Neuroimaging technologies and more recent research, how-

ever, have shown that certain pleasurable activities, such as gambling, shopping, and sex, can also co-opt the brain. Although the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)* describes multiple addictions, each tied to a specific substance or activity, consensus is emerging that these may represent multiple expressions of a common underlying brain process.

### From liking to wanting

Nobody starts out intending to develop an addiction, but many people get caught in its snare. According to the latest government statistics, nearly 23 million Americans—almost one in 10—are addicted to alcohol or other drugs. More than two-thirds of people with addiction abuse alcohol. The top three drugs causing addiction are marijuana, opioid (narcotic) pain relievers, and cocaine.

Genetic vulnerability contributes to the risk of developing an addiction. Twin and adoption studies show that about 40% to 60% of susceptibility to addiction is hereditary. But behavior plays a key role, especially when it comes to reinforcing a habit.

**Pleasure principle.** The brain registers all pleasures in the same way, whether they originate with a psychoactive drug, a monetary reward, a sexual encounter, or a satisfying meal. In the brain, pleasure has a distinct signature: the release of the neurotransmitter dopamine in the nucleus accumbens, a cluster of nerve cells lying underneath the cerebral cortex (see illustration, page 3). Dopamine release in the nucleus accumbens is so consistently tied with pleasure that neuroscientists refer to the region as the brain's pleasure center.

All drugs of abuse, from nicotine to heroin, cause a particularly powerful surge. ▶

## INSIDE

When depression starts in the neck. . . . . 4  
Treating an underactive thyroid gland may improve mood.

Expressive writing for mental health. . . . . 6  
Putting an experience into words may ease stress and trauma.

In brief. . . . . 7  
Long-term results of deep brain stimulation for depression; More evidence that exercise aids the brain.

Commentary. . . . . 8  
FDA panel finds no link between food colorings and hyperactivity in most children.

## WHAT'S NEW

### Special Health Report

**Hands**  
*Strategies for strong, pain-free hands*

To order, call 877-649-9457 (toll-free) or visit us online at [www.health.harvard.edu](http://www.health.harvard.edu).



Receive Harvard Health Publications' free e-mail newsletter

Go to [www.health.harvard.edu](http://www.health.harvard.edu) to subscribe to *HEALTHbeat*. This free, weekly e-mail newsletter brings you health tips, advice, and information on a wide range of topics.

You can also join in discussion with experts from Harvard Health Publications and folks like you on various health topics, medical news, and views by reading the Harvard Health Blog ([www.health.harvard.edu/blog](http://www.health.harvard.edu/blog)).

## CONTACT US

Write to us at [mental\\_letter@hms.harvard.edu](mailto:mental_letter@hms.harvard.edu)

For customer service, write us at [harvardMH@strategicfulfillment.com](mailto:harvardMH@strategicfulfillment.com)



# Harvard Mental Health Letter



Editorial Board members are associated with Harvard Medical School and affiliated institutions. They review all published articles.

**Editor in Chief** Michael Craig Miller, MD  
**Editor** Ann MacDonald  
**Contributor** Katherine Perch  
**Founding Editor** Lester Grinspoon, MD  
**Editorial Board** Mary Anne Badaracco, MD  
Jonathan F. Borus, MD  
Christopher B. Daly  
Sandra Dejong, MD  
Frank W. Drislane, MD  
Anne K. Fishel, PhD  
Donald C. Goff, MD  
Stuart Goldman, MD  
Alan I. Green, MD  
Shelly Greenfield, MD, MPH  
Thomas G. Gutheil, MD  
Michael Hirsch, MD  
Matcheri S. Keshavan, MD  
Kimberlyn Leary, PhD, ABPP  
Robert W. McCarley, MD  
Michael J. Mufson, MD  
Andrew A. Nierenberg, MD  
Scott L. Rauch, MD, PhD  
Hester H. Schnipper, LICSW, BCD  
Janna M. Smith, LICSW, BCD  
Caroline L. Watts, EdD  
Barbara Wolfe, PhD, RN

**Art Director** Heather Derocher  
**Production Editor** Nicole Wall  
**Illustrator** Scott Leighton

**Customer Service**  
**Phone** 877-649-9457 (toll-free)  
**E-mail** harvardMH@strategicfulfillment.com  
**Online** www.health.harvard.edu/customer\_service  
**Letters** Harvard Mental Health Letter  
P.O. Box 9308, Big Sandy, TX 75755-9308

**Subscriptions** \$72 per year (U.S.)  
**Bulk Subscriptions**  
StayWell Consumer Health Publishing  
1 Atlantic St., Suite 604, Stamford, CT 06901  
888-456-1222, ext. 31106 (toll-free)  
203-653-6266  
ddewitt@staywell.com

**Corporate Sales and Licensing**  
StayWell Consumer Health Publishing  
1 Atlantic St., Suite 604, Stamford, CT 06901  
jmittchell@staywell.com

**Editorial Correspondence**  
**E-mail** mental\_letter@hms.harvard.edu  
**Letters** Harvard Mental Health Letter  
10 Shattuck St., 2nd Floor, Boston, MA 02115

**Permissions**  
Copyright Clearance Center, Inc.  
**Online** www.copyright.com

Published by Harvard Health Publications,  
a division of Harvard Medical School

**Editor in Chief** Anthony L. Komaroff, MD  
**Publishing Director** Edward Coburn

© 2011 Harvard University (ISSN 0884-3783)  
Proceeds support the research efforts of Harvard Medical School.

Harvard Health Publications  
10 Shattuck St., 2nd Floor, Boston, MA 02115

The goal of the Harvard Mental Health Letter is to interpret timely mental health information. Its contents are not intended to provide advice for individual problems. Such advice should be offered only by a person familiar with the detailed circumstances in which the problem arises. We are interested in comments and suggestions about the content; unfortunately, we cannot respond to all inquiries.

PUBLICATIONS MAIL AGREEMENT NO. 40906010  
RETURN UNDELIVERABLE CANADIAN ADDRESSES TO:  
CIRCULATION DEPT., 1415 JANETTE AVENUE,  
WINDSOR, ON N8X 1Z1 • E-mail: ddewitt@staywell.com

## Addiction continued

of dopamine in the nucleus accumbens. The likelihood that the use of a drug or participation in a rewarding activity will lead to addiction is directly linked to the speed with which it promotes dopamine release, the intensity of that release, and the reliability of that release. Even taking the same drug through different methods of administration can influence how likely it is to lead to addiction. Smoking a drug or injecting it intravenously, as opposed to swallowing it as a pill, for example, generally produces a faster, stronger dopamine signal and is more likely to lead to drug misuse.

### Learning process.

Scientists once believed that the experience of pleasure alone was enough to prompt people to continue seeking an addictive substance or activity. But more recent research suggests that the situation is more complicated. Dopamine not only contributes to the experience of pleasure, but also plays a role in learning and memory—two key elements in the transition from liking something to becoming addicted to it.

According to the current theory about addiction, dopamine interacts with another neurotransmitter, glutamate, to take over the brain's system of reward-related learning. This system has an important role in sustaining life because it links activities needed for human survival (such as eating and sex) with pleasure and reward. The reward circuit in the brain includes areas involved with motivation and memory as well as with pleasure. Addictive substances and behaviors stimulate the same circuit—and then overload it.

Repeated exposure to an addictive

substance or behavior causes nerve cells in the nucleus accumbens and the prefrontal cortex (the area of the brain involved in planning and executing tasks) to communicate in a way that couples liking something with wanting it, in turn driving us to go after it. That is, this process motivates us to take action to seek out the source of pleasure.

Tolerance and compulsion. Over time, the brain adapts in a way that actually makes the sought-after substance or activity less pleasurable.

In nature, rewards usually come only with time and effort. Addictive drugs and behaviors provide a shortcut, flooding the brain with dopamine and other neurotransmitters. Our brains do not have an easy way to withstand the onslaught.

Addictive drugs, for example, can release two to 10 times the

amount of dopamine that natural rewards do, and they do it more quickly and more reliably. In a person who becomes addicted, brain receptors become overwhelmed. The brain responds by producing less dopamine or eliminating dopamine receptors—an adaptation similar to turning the volume down on a loudspeaker when noise becomes too loud.

As a result of these adaptations, dopamine has less impact on the brain's reward center. People who develop an addiction typically find that, in time, the desired substance no longer gives them as much pleasure. They have to take more of it to obtain the same dopamine "high" because their brains have adapted—an effect known as tolerance.

At this point, compulsion takes over. The pleasure associated with an addic-

## Resources

National Clearinghouse for Alcohol and Drug Information  
P.O. Box 2345  
Rockville, MD 20847  
800-729-6686 (toll-free)  
<http://ncadi.samhsa.gov>

National Institute on Alcohol Abuse and Alcoholism  
5635 Fishers Lane, MSC 9304  
Bethesda, MD 20892  
301-443-3860  
[www.niaaa.nih.gov](http://www.niaaa.nih.gov)

National Institute on Drug Abuse  
6001 Executive Blvd., Room 5213  
Bethesda, MD 20892  
301-443-1124  
[www.nida.nih.gov](http://www.nida.nih.gov)

Substance Abuse and Mental Health Services Administration  
1 Choke Cherry Road  
Rockville, MD 20857  
877-276-4727 (toll-free)  
[www.samhsa.gov](http://www.samhsa.gov)



tive drug or behavior subsides—and yet the memory of the desired effect and the need to recreate it (the *wanting*) persists. It's as though the normal machinery of motivation is no longer functioning.

The learning process mentioned earlier also comes into play. The hippocampus and the amygdala store information about environmental cues associated with the desired substance, so that it can be located again. These memories help create a conditioned response—intense craving—whenever the person encounters those environmental cues.

Cravings contribute not only to addiction but to relapse after a hard-won sobriety. A person addicted to heroin may be in danger of relapse when he sees a hypodermic needle, for example, while another person might start to drink again after seeing a bottle of whiskey. Conditioned learning helps explain why people who develop an addiction risk relapse even after years of abstinence.


### The long road to recovery

Because addiction is learned and stored in the brain as memory, recovery is a slow and hesitant process in which the influence of those memories diminishes.

About 40% to 60% of people with a drug addiction experience at least one relapse after an initial recovery. While this may seem discouraging, the relapse rate is similar to that in other chronic diseases, such as high blood pressure and asthma, where 50% to 70% of people each year experience a recurrence of symptoms significant enough to require medical intervention.

Fortunately a number of effective treatments exist for addiction, usually combining self-help strategies, psychotherapy, and rehabilitation. For some types of addictions, medication may also help. (We've covered specific addictions in detail in previous articles. See *Harvard Mental Health Letter*,

### The brain's reward center



Addictive drugs provide a shortcut to the brain's reward system by flooding the nucleus accumbens with dopamine. The hippocampus lays down memories of this rapid sense of satisfaction, and the amygdala creates a conditioned response to certain stimuli.

March 2009, January 2010, and April 2011.)

The precise plan varies based on the nature of the addiction, but all treatments are aimed at helping people to unlearn their addictions while adopting healthier coping strategies—truly a brain-based recovery program. ♥

Benowitz NL. "Nicotine Addiction," *The New England Journal of Medicine* (June 17, 2010): Vol. 362, No. 24, pp. 2295–303.

Brady KT, et al., eds. *Women and Addiction: A Comprehensive Handbook* (The Guilford Press, 2009).

Chandler RK, et al. "Treating Drug Abuse and Addiction in the Criminal Justice System: Improving Public Health and Safety," *Journal of the American Medical Association* (Jan. 14, 2009): Vol. 301, No. 2, pp. 183–90.

Greenfield SE, et al. "Substance Abuse Treatment Entry, Retention, and Outcome in Women: A Review of the Literature," *Drug and Alcohol Dependence* (Jan. 5, 2007): Vol. 86, No. 1, pp. 1–21.

Koob GF, et al. "Neurocircuitry of Addiction," *Neuropsychopharmacology* (Jan. 2010): Vol. 35, No. 1, pp. 217–38.

McLellan AT, et al. "Drug Dependence, A Chronic Medical Illness: Implications for Treatment, Insurance, and Outcomes Evaluation," *Journal of the American Medical Association*

(Oct. 4, 2000): Vol. 284, No. 13, pp. 1689–95.

National Institute on Drug Abuse. *Drugs, Brains, and Behavior: The Science of Addiction* (National Institutes of Health, Aug. 2010).

Polosa R, et al. "Treatment of Nicotine Addiction: Present Therapeutic Options and Pipeline Developments," *Trends in Pharmacological Sciences* (Jan. 20, 2011): E-publication.

Potenza MN, et al. "Neuroscience of Behavioral and Pharmacological Treatments for Addictions," *Neuron* (Feb. 24, 2011): Vol. 69, No. 4, pp. 695–712.

Shaffer HJ, et al. "Toward a Syndrome Model of Addiction: Multiple Expressions, Common Etiology," *Harvard Review of Psychiatry* (Nov.–Dec. 2004): Vol. 12, No. 6, pp. 367–74. \*

Stead LF, et al. "Nicotine Replacement Therapy for Smoking Cessation," *Cochrane Database of Systematic Reviews* (Jan. 23, 2008): Doc. No. CD000146.

Substance Abuse and Mental Health Services Administration. *National Survey on Drug Use & Health*, 2009.

\* Harvard Mental Health Letter subscribers can obtain a special discounted subscription to the *Harvard Review of Psychiatry* by visiting the Web site <http://informaworld.com/hrp> or by calling 212-520-2763.

For more references, please see [www.health.harvard.edu/mentalextra](http://www.health.harvard.edu/mentalextra).